

# UNCLASSIFIED

AD NUMBER
AD842585
NEW LIMITATION CHANGE
TO Approved for public release, distribution unlimited
FROM Distribution authorized to U.S. Gov't. agencies and their contractors; Critical Technology; OCT 1968. Other requests shall be referred to Commanding Officer, Edgewood Arsenal, Attn: SMUEA-TSTI-T, Edgewood Arsenal, MD 21010.
AUTHORITY
USAEA ltr, 22 Dec 1971

THIS PAGE IS UNCLASSIFIED

AD 842585

AD

**EDGEWOOD ARSENAL  
TECHNICAL REPORT**



**EATR 4231**

**METAL ION COMPLEXATION OF MODEL IMIDES**

by

Raymond A. Mackay  
Edward J. Poziomek

October 1968



**DEPARTMENT OF THE ARMY  
EDGEWOOD ARSENAL  
Research Laboratories  
Physical Research Laboratory  
Edgewood Arsenal, Maryland 21010**

ADD	NO. 1	NO. 2	NO. 3	NO. 4	NO. 5	NO. 6	NO. 7	NO. 8	NO. 9	NO. 10
1	2	3	4	5	6	7	8	9	10	11
12	13	14	15	16	17	18	19	20	21	22
23	24	25	26	27	28	29	30	31	32	33
34	35	36	37	38	39	40	41	42	43	44
45	46	47	48	49	50	51	52	53	54	55
56	57	58	59	60	61	62	63	64	65	66
67	68	69	70	71	72	73	74	75	76	77
78	79	80	81	82	83	84	85	86	87	88
89	90	91	92	93	94	95	96	97	98	99
100	101	102	103	104	105	106	107	108	109	110
111	112	113	114	115	116	117	118	119	120	121
122	123	124	125	126	127	128	129	130	131	132
133	134	135	136	137	138	139	140	141	142	143
144	145	146	147	148	149	150	151	152	153	154
155	156	157	158	159	160	161	162	163	164	165
166	167	168	169	170	171	172	173	174	175	176
177	178	179	180	181	182	183	184	185	186	187
188	189	190	191	192	193	194	195	196	197	198
199	200	201	202	203	204	205	206	207	208	209
210	211	212	213	214	215	216	217	218	219	220
221	222	223	224	225	226	227	228	229	230	231
232	233	234	235	236	237	238	239	240	241	242
243	244	245	246	247	248	249	250	251	252	253
254	255	256	257	258	259	260	261	262	263	264
265	266	267	268	269	270	271	272	273	274	275
276	277	278	279	280	281	282	283	284	285	286
287	288	289	290	291	292	293	294	295	296	297
298	299	300	301	302	303	304	305	306	307	308
309	310	311	312	313	314	315	316	317	318	319
320	321	322	323	324	325	326	327	328	329	330
331	332	333	334	335	336	337	338	339	340	341
342	343	344	345	346	347	348	349	350	351	352
353	354	355	356	357	358	359	360	361	362	363
364	365	366	367	368	369	370	371	372	373	374
375	376	377	378	379	380	381	382	383	384	385
386	387	388	389	390	391	392	393	394	395	396
397	398	399	400	401	402	403	404	405	406	407
408	409	410	411	412	413	414	415	416	417	418
419	420	421	422	423	424	425	426	427	428	429
430	431	432	433	434	435	436	437	438	439	440
441	442	443	444	445	446	447	448	449	450	451
452	453	454	455	456	457	458	459	460	461	462
463	464	465	466	467	468	469	470	471	472	473
474	475	476	477	478	479	480	481	482	483	484
485	486	487	488	489	490	491	492	493	494	495
496	497	498	499	500	501	502	503	504	505	506
507	508	509	510	511	512	513	514	515	516	517
518	519	520	521	522	523	524	525	526	527	528
529	530	531	532	533	534	535	536	537	538	539
540	541	542	543	544	545	546	547	548	549	550
551	552	553	554	555	556	557	558	559	560	561
562	563	564	565	566	567	568	569	570	571	572
573	574	575	576	577	578	579	580	581	582	583
584	585	586	587	588	589	590	591	592	593	594
595	596	597	598	599	600	601	602	603	604	605
606	607	608	609	610	611	612	613	614	615	616
617	618	619	620	621	622	623	624	625	626	627
628	629	630	631	632	633	634	635	636	637	638
639	640	641	642	643	644	645	646	647	648	649
650	651	652	653	654	655	656	657	658	659	660
661	662	663	664	665	666	667	668	669	670	671
672	673	674	675	676	677	678	679	680	681	682
683	684	685	686	687	688	689	690	691	692	693
694	695	696	697	698	699	700	701	702	703	704
705	706	707	708	709	710	711	712	713	714	715
716	717	718	719	720	721	722	723	724	725	726
727	728	729	730	731	732	733	734	735	736	737
738	739	740	741	742	743	744	745	746	747	748
749	750	751	752	753	754	755	756	757	758	759
760	761	762	763	764	765	766	767	768	769	770
771	772	773	774	775	776	777	778	779	780	781
782	783	784	785	786	787	788	789	790	791	792
793	794	795	796	797	798	799	800	801	802	803
804	805	806	807	808	809	810	811	812	813	814
815	816	817	818	819	820	821	822	823	824	825
826	827	828	829	830	831	832	833	834	835	836
837	838	839	840	841	842	843	844	845	846	847
848	849	850	851	852	853	854	855	856	857	858
859	860	861	862	863	864	865	866	867	868	869
870	871	872	873	874	875	876	877	878	879	880
881	882	883	884	885	886	887	888	889	890	891
892	893	894	895	896	897	898	899	900	901	902
903	904	905	906	907	908	909	910	911	912	913
914	915	916	917	918	919	920	921	922	923	924
925	926	927	928	929	930	931	932	933	934	935
936	937	938	939	940	941	942	943	944	945	946
947	948	949	950	951	952	953	954	955	956	957
958	959	960	961	962	963	964	965	966	967	968
969	970	971	972	973	974	975	976	977	978	979
980	981	982	983	984	985	986	987	988	989	990
991	992	993	994	995	996	997	998	999	1000	1001

### Distribution Statement

This document is subject to special export controls and each transmittal to foreign governments or foreign nationals may be made only with prior approval of the Commanding Officer, Edgewood Arsenal, ATTN: SMUEA-TSTI-T, Edgewood Arsenal, Maryland 21010.

### Disclaimer

The findings in this report are not to be construed as an official Department of the Army position unless so designated by other authorized documents.

### Disposition

When this report is no longer needed, destroy it. Do not return it to the originator.

EDGEWOOD ARSENAL TECHNICAL REPORT

EATR 4231

METAL ION COMPLEXATION OF MODEL IMIDES

by

Raymond A. Mackay  
Edward J. Poziomek

Defensive Research Department

October 1968

This document is subject to special export controls and each transmittal to foreign governments or foreign nationals may be made only with prior approval of the Commanding Officer, Edgewood Arsenal, ATTN: SMUEA-TSTI-T, Edgewood Arsenal, Maryland 21010.

Task 1B622401A10204

DEPARTMENT OF THE ARMY  
EDGEWOOD ARSENAL  
Research Laboratories  
Physical Research Laboratory  
Edgewood Arsenal, Maryland 21010

### FOREWORD

The work described in this report was authorized under Task 1B622401A10204, Detection and Warning Investigations (U). The work was started in October 1967 and completed in April 1968. The experimental data are contained in notebooks 7798 and 7908.

Reproduction of this document in whole or in part is prohibited except with permission of the Commanding Officer, Edgewood Arsenal, ATTN: SMUEA-TSTI-T, Edgewood Arsenal, Maryland 21010; however, Defense Documentation Center is authorized to reproduce the document for United States Government purposes.

The information in this document has not been cleared for release to the general public.

### Acknowledgments

Elemental analyses were performed by the Analytical Chemistry Department.

### DIGEST

The purpose of this investigation was to study metal ion complexation of model imides. This was accomplished by observing the effect of metal ion complexation on the carbonyl stretching frequencies of two imides, N-cyclohexyl-N-formylacetamide and 3,5-dioxopyrrolizidine. Two metal complexes were isolated,  $(\text{CuCl}_2)_3(\text{N-cyclohexyl-N-formylacetamide})_2$  and  $\text{Co}(3,5\text{-dioxopyrrolizidine})_2(\text{ClO}_4)_2$ .

It is concluded on the basis of analyzing the infrared absorption data that when an imide is fixed in a conformation represented by the one for 3,5-dioxopyrrolizidine then chelate formation takes place readily. However, when free rotation around the imide C-N bonds is possible then metal complexation takes place only at one carbonyl and then at the oxygen with the highest electron density.

## CONTENTS


	<u>Page</u>
I. INTRODUCTION . . . . .	7
II. EXPERIMENTATION . . . . .	8
A. Preparation of Compounds . . . . .	8
B. Infrared Spectra . . . . .	9
III. DISCUSSION . . . . .	9
IV. CONCLUSIONS . . . . .	11
LITERATURE CITED . . . . .	15
DISTRIBUTION LIST . . . . .	17

## METAL ION COMPLEXATION OF MODEL IMIDES


### I. INTRODUCTION.

Because of current interest by the Detection Research Branch in the nature of intermediates found in metal ion catalyzed reactions of amides with electrophiles, we undertook a study of the metal ion complexation of imides. Imides are formed as a result of allowing formamides to react with acylating agents.

The effect of metal ion complexation on the infrared absorption spectra of amides has been thoroughly investigated.<sup>1-4</sup> The spectra of metal ion-imide complexes have also been recently examined.<sup>5-10</sup> These studies all involved acyclic symmetrical imides (e. g., diacetamides), and we do not believe that such studies have been reported using acyclic unsymmetrical or cyclic imides. The interesting feature is that several metal complexes are possible since four configurations (A-D) can usually be written. The only



conformation which is capable of chelate formation with a metal ion is (A). The other conformations should coordinate at only one of the oxygens. (The nitrogen atom is also capable of complexing, but this does not occur for first-row transition metals.) The previous studies have shown that imides can act as both mono- and bidentate ligands, with coordination to the metal via oxygen. In the majority of cases, the imide was bidentate, complexation apparently stabilizing the conformation A. In the cases where the imide was monodentate, both oxygens were usually coordinated but to different metal atoms. We chose to examine the effect of metal ion complexation on the carbonyl stretching frequency of two imides, N-cyclohexyl-N-formylacetamide (I) and 3,5-dioxopyrrolizidine (II). Compound II has configuration A.



while (I) may possess any or all of the forms shown.



## II. EXPERIMENTATION.

### A. Preparation of Compounds.

#### 1. N-Alkyl-N-formylacetamides.

These compounds were prepared by Hoy and Poziomek.<sup>11</sup>

#### 2. 3, 5-Dioxopyrrolizidine.

A sample of this compound was obtained through the courtesy of Dr. Arnold J. Gordon, Catholic University of America. The particular sample was prepared by Michael and Flitech.<sup>12</sup>

#### 3. (CuCl<sub>2</sub>)<sub>3</sub>(N-Cyclohexyl-N-formylacetamide)<sub>2</sub>.

CuCl<sub>2</sub>·2H<sub>2</sub>O (1.7 grams) was dehydrated by stirring for 1 hour with 10 ml of ethyl orthoformate. The salt did not dissolve but formed an olive green slurry. Upon addition of 7 ml of N-cyclohexyl-N-formylacetamide, the slurry began to form a mustard colored polycrystalline solid. After stirring for 1 hour to insure complete reaction, the solid was filtered off, washed well with ether, and dried in a vacuum desiccator over P<sub>4</sub>O<sub>10</sub>, mp 192° to 193° decomposes.

Anal. Calcd. for Cu<sub>3</sub>Cl<sub>6</sub>C<sub>18</sub>H<sub>30</sub>N<sub>2</sub>O<sub>4</sub>: Cu, 25.6; Cl, 28.8; C, 29.2; H, 4.0; N, 3.8. Found: Cu, 25.3; Cl, 28.7; C, 28.3; H, 4.1; N, 4.0.

#### 4. Co(3, 5-Dioxopyrrolizidine)<sub>2</sub>(ClO<sub>4</sub>)<sub>2</sub>·H<sub>2</sub>O.

Co(ClO<sub>4</sub>)<sub>2</sub>·6H<sub>2</sub>O (0.85 gram) was dehydrated by stirring with 6 ml of ethyl orthoformate and 0.85 gram of 3, 5-dioxopyrrolizidine was then added. Stirring was allowed to continue for 1 hour. The light pink solid obtained was filtered off, washed with a small amount of ethanol, then with ether, and dried in a vacuum desiccator over P<sub>4</sub>O<sub>10</sub>, mp 272° to 276° decomposes.

Anal. Calcd. for CoCl<sub>2</sub>C<sub>14</sub>H<sub>20</sub>N<sub>2</sub>O<sub>13</sub>: Co, 10.6; Cl 12.8; C, 30.3; H, 3.6; N, 5.1. Found: Co, 10.6; Cl, 12.4; C, 31.0; H, 3.8; N, 5.1.

### B. Infrared Spectra.

A Perkin-Elmer model 521 grating spectrophotometer was employed to record the infrared spectra. The samples were prepared as neat liquids or Nujol mulls held between KRS-5 (TlI/Br) plates. The carbonyl stretching region was calibrated with polystyrene. The data are given in the table. Results for some N-alkyl amides are also included for purposes of comparison.

Table. Infrared Absorption Data

Compound	Carbonyl stretching frequency $\text{cm}^{-1}$ <sup>a/</sup>
n-Butylformamide <sup>b/</sup>	1670
Sec-butylformamide <sup>b/</sup>	1670
Cyclohexylformamide <sup>b/</sup>	1665
N-(n-butyl)acetamide <sup>c/</sup>	1645
N-(n-butyl)-N-formylacetamide <sup>b/</sup>	1720, 1670
N-(sec-butyl)-N-formylacetamide <sup>b/</sup>	1730, 1687
N-cyclohexyl-N-formylacetamide <sup>b/</sup>	1720, 1672
(CuCl <sub>2</sub> ) <sub>3</sub> (N-cyclohexyl-N-formylacetamide) <sub>2</sub> <sup>c/</sup>	1741, 1618
3, 5-Dioxopyrrolizidine <sup>c/</sup>	1770, 1688
Co(3, 5-dioxopyrrolizidine) <sub>2</sub> (ClO <sub>4</sub> ) <sub>2</sub> ·H <sub>2</sub> O <sup>c/</sup>	1733, 1642

a/ Errors are ca  $\pm 3 \text{ cm}^{-1}$ .


b/ Neat liquid.

c/ Mull.


### III. DISCUSSION.

The carbonyl stretching frequencies of N-substituted amides (III) are not appreciably affected by changes in R, when R is alkyl. This

also holds true for imides of the N-alkyl-N-formylacetamide type (IV).



III




IV

However, the carbonyl frequency is influenced by whether R' is H or alkyl. For example,  $\gamma$  C=O occurs at  $1670 \text{ cm}^{-1}$  for n-butylformamide (R'=H) and  $1645 \text{ cm}^{-1}$  for N-(n-butyl)acetamide (R'=CH<sub>3</sub>) (table). Both carbonyl stretching bands observed in spectra of N-alkyl-N-formylacetamides appear at higher frequencies than in either alkylformamides or acetamides. This is because the pair of electrons from nitrogen are delocalized over two carbonyl groups (structures V and VI) in comparison to one (structure VII) in the alkyl amide.



V

VI



VII

Of the pair of carbonyl absorption bands found for N-alkyl-N-formylacetamides, the higher frequency one is assigned to the formyl carbonyl. The distinction between formyl and acetyl is very evident by comparing N-(n-butyl)-N-formylacetamide with n-butylformamide and N-(n-butyl)acetamide (table). The acetyl carbonyl absorbs at lower frequencies because of the inductive and hyperconjugative effects from the methyl group. The carbonyl absorption of 3, 5-dioxopyrrolizidine is at higher frequency than in the N-alkyl-N-formylacetamides because of ring strain.\* It may be noted that while both carbonyl groups are stereochemically equivalent in 3, 5-dioxopyrrolizidine, there are two carbonyl bands. This is due to the strong coupling of the two carbonyl vibrational modes.

\* It is known that lactam carbonyl absorption is normal in unstrained rings of six carbon atoms, but shifts toward higher frequencies in smaller rings because of ring strain.<sup>13</sup>


For the  $(\text{CuCl}_2)_3(\text{N-cyclohexyl-N-formylacetamide})_2$  complex, \* the spectrum (figure 1) shows clearly that only the acetyl groups is complexed. The formyl band at  $1720\text{ cm}^{-1}$  in the free ligand shifts to somewhat higher frequency ( $1741\text{ cm}^{-1}$ ), while the  $1672\text{ cm}^{-1}$  acetyl band drops to  $1618\text{ cm}^{-1}$ . Preferential coordination to the acetyl oxygen was expected since the oxygen electron density relative to formyl is higher (methyl inductive and hyperconjugative effects). In the case of  $\text{Co}(3,5\text{-dioxopyrrolizidine})_2(\text{ClO}_4)_2$  (figure 2), both carbonyl oxygens are bound to the metal as evidenced by the shifts of both carbonyl bands to lower frequency. The absence of chelate formation using  $\text{CuCl}_2$  and N-cyclohexyl-N-formylacetamide is understandable since electrostatic repulsion between the oxygen atoms would disfavor configuration A. Complexation of N-cyclohexyl-N-formylacetamide with  $\text{Co}(\text{ClO}_4)_2$  gave a product that could not be purified but whose infrared absorption spectrum also indicated the presence of unchelated carbonyl.

Our studies using two model imides (I and II) appear to indicate that when an imide is fixed in conformation A, chelate formation takes place readily. However, when free rotation around the C-N bonds is possible then metal complexation takes place only at one carbonyl and then at the oxygen with the highest electron density.

#### IV. CONCLUSIONS.

It is concluded on the basis of analyzing the infrared absorption data that when an imide is fixed in a conformation represented by the one for 3,5-dioxopyrrolizidine then chelate formation takes place readily. However, when free rotation around the imide C-N bonds is possible then metal complexation takes place only at one carbonyl and then at the oxygen with the highest electron density.

- \* A knowledge of the structure of this complex is not necessary for the purpose of this work. However, the electronic spectrum of the solid indicates a planar or tetragonal configuration about copper. Other copper compounds of the stoichiometry  $(\text{CuCl}_2)_3\text{L}_2$  are known, which generally have the structure, where each copper atom is in a square planar



configuration with a weak tetragonal interaction because of a chlorine atom from another  $(\text{CuCl}_2)_3\text{L}_2$  monomer unit in the layers above and below, 14, 15




Figure 1. Infrared Absorption Spectra of  $(\text{CuCl}_2)_3(\text{N-cyclohexyl-N-formylacetamide})_2$  (----) and N-cyclohexyl-N-formylacetamide (—).



Figure 2. Infrared Absorption Spectra of  $\text{Co}(3, 5\text{-dioxopyrrolizidine})_2 (\text{ClO}_4)_2 \cdot \text{H}_2\text{O}$  (—) and 3, 5-Dioxopyrrolizidine (----).

### LITERATURE CITED

1. Bull, W. E., Madan, S. K., and Willis, J. E. Amides as Ligands. I. Metallic Complexes of N,N-Dimethylacetamide. *Inorg. Chem.* 2, 303 (1963).
2. Drago, R. S., Meek, D. W., Joesten, M. D., and La Roche, L. Spectrochemical Studies of a Series of Amides as Ligands With Nickel (II) and Chromium (III). *Inorg. Chem.* 2, 124 (1963).
3. Madan, S. K. Amides as Ligands. VII. Complexes of N-Methyl- $\gamma$ -butyrolactam With Nontransition Metals, *Inorg. Chem.* 6, 421 (1967).
4. Madan, S. K., and Sturr, J. A. Amides as Ligands. VIII. Inorganic Complexes of  $\gamma$ -Butyrolactam and N-Methyl- $\gamma$ -butyrolactam. *J. Inorg. Nucl. Chem.* 29, 1669 (1967).
5. Gentile, P. S., and Shankoff, T. A. Solid-Solid Interactions. III. Complexes of Diacetamide and Alkali and Alkaline Earth Metal Salts. *J. Inorg. Nucl. Chem.* 27, 2301 (1965).
6. Gentile, P. S. and Shankoff, T. A. Complexes of Diacetamide and Divalent Transition Metal Salts. *J. Inorg. Nucl. Chem.* 28, 1283 (1966).
7. Kraihanzel, C. S., and Grenda, S. C. Acyclic Imides as Ligands. I. Diacetamide Complexes of Manganese (II), Iron (II), Cobalt (II), Nickel (II), Copper (II), and Zinc (II) Perchlorates. *Inorg. Chem.* 4, 1037 (1965).
8. Gentile, P. S., Shankoff, T. A., and Carlotto, J. Complexes and Configurational Isomers of Diacetamide with Group IIB Metal Salts. *J. Inorg. Nucl. Chem.* 28, 979 (1966).
9. Gentile, P. S., Shankoff, T. A. Polymerization Isomers of Copper (II) Chloride. *J. Inorg. Nucl. Chem.* 28, 1125 (1966).
10. Jain, S. C., and Rivest, R. Coordination Complexes of Group (IV) Halides. *J. Inorg. Nucl. Chem.* 29, 2787 (1967).
11. Hoy, D. J., and Poziomek, E. J. Studies in the Pyrolysis of N-Formylacetamides. The Imide-Isimide Rearrangement. *J. Org. Chem.* (November 1968). (In press).

12. Micheel, F., and Flitsch, W. Eine einfache Synthese des 3,4-Dioxo-pyrrolizidins (pyrrolizidinderivate III). Ber. 88, 509 (1955).

13. Bellamy, L. J. The Infrared Spectra of Complex Molecules. p 213. John Wiley and Sons, Inc. 1958.

14. Willett, R. D., and Rundle, R. E. Crystal Structure of  $\text{Cu}_2\text{Cl}_4(\text{CH}_3\text{CN})_2$ ,  $\text{Cu}_3\text{Cl}_6(\text{CH}_3\text{CN})_2$ , and  $\text{Cu}_5\text{Cl}_{10}(\text{C}_3\text{H}_7\text{OH})_2$ . J. Chem. Phys. 40, 838 (1964).

15. Whyman, R., Copley, D. B., and Hatfield, W. E. Spin-Spin Coupling in Binuclear Complexes. IV. Substituted Heterocyclic N-Oxide Complexes of Copper (II) Halides. III. Magnetic and Spectral Properties of Complexes With Substituted Quinoline N-Oxides. J. Amer. Chem. Soc. 89, 3135 (1967).



UNCLASSIFIED

Security Classification

DOCUMENT CONTROL DATA - R & D

(Security classification of title, body of abstract and indexing annotation must be entered when the overall report is classified)

1. ORIGINATING ACTIVITY (Corporate author) CO, Edgewood Arsenal ATTN: SMUEA-RPRE Edgewood Arsenal, Maryland 21010		2a. REPORT SECURITY CLASSIFICATION UNCLASSIFIED													
3. REPORT TITLE METAL ION COMPLEXATION OF MODEL IMIDES		2b. GROUP NA													
4. DESCRIPTIVE NOTES (Type of report and inclusive dates) This work was started in October 1967 and completed in April 1968.															
5. AUTHOR(S) (First name, middle initial, last name) Mackay, Raymond A., and Poziomek, Edward J.															
6. REPORT DATE October 1968		7a. TOTAL NO. OF PAGES 21	7b. NO. OF REFS 15												
8. CONTRACT OR GRANT NO.		9a. ORIGINATOR'S REPORT NUMBER(S) EATR 4231													
b. PROJECT NO.		9b. OTHER REPORT HO(S) (Any other numbers that may be assigned this report) NA													
c. Task No. 1B622401A10204															
10. DISTRIBUTION STATEMENT This document is subject to special export controls and each transmittal to foreign governments or foreign nationals may be made only with prior approval of the Commanding Officer, Edgewood Arsenal, ATTN: SMUEA-TSTI-T, Edgewood Arsenal, Maryland 21010.															
11. SUPPLEMENTARY NOTES Detection and warning investigations		12. SPONSORING MILITARY ACTIVITY NA													
13. ABSTRACT (U) - The purpose of this investigation was to study metal ion complexation of model imides. This was accomplished by observing the effect of metal ion complexation on the carbonyl stretching frequencies of two imides, N-cyclohexyl-N-formylacetamide and 3,5-dioxopyrrolizidine. Two metal complexes were isolated, $(\text{CuCl}_2)_3(\text{N-cyclohexyl-N-formylacetamide})_2$ and $\text{Co}(3,5\text{-dioxopyrrolizidine})_2(\text{C}_{10}\text{H}_{16}\text{O}_2)_2$ . It is concluded on the basis of analyzing the infrared absorption data that when an imide is fixed in a conformation represented by the one for 3,5-dioxopyrrolizidine then chelate formation takes place readily. However, when free rotation around the imide C-N bonds is possible then metal complexation takes place only at one carbonyl and then at the oxygen with the highest electron density.															
14. KEYWORDS <table border="0"><tr><td>Metal ion complexation</td><td>Acetamide</td></tr><tr><td>3,5-Dioxopyrrolizidine</td><td>Detection</td></tr><tr><td>Cobalt perchlorate</td><td>Imides</td></tr><tr><td>Copper chloride</td><td>Infrared</td></tr><tr><td>Formamides</td><td>Chelate</td></tr><tr><td>Carbonyl</td><td>N-Cyclohexyl-N-formylacetamide</td></tr></table>				Metal ion complexation	Acetamide	3,5-Dioxopyrrolizidine	Detection	Cobalt perchlorate	Imides	Copper chloride	Infrared	Formamides	Chelate	Carbonyl	N-Cyclohexyl-N-formylacetamide
Metal ion complexation	Acetamide														
3,5-Dioxopyrrolizidine	Detection														
Cobalt perchlorate	Imides														
Copper chloride	Infrared														
Formamides	Chelate														
Carbonyl	N-Cyclohexyl-N-formylacetamide														

DD FORM 1473

REPLACES DD FORM 1473, 1 JAN 64, WHICH IS OBSOLETE FOR ARMY USE.

UNCLASSIFIED

Security Classification